Fish are the most Commonand dvuse group of Vertebratiganisms in the ocean
Their origins date back 500 million years
There are about 32,000 different species of fish WOY du I de la both Fresh and Salt water)
Fish are classified as Chordato meaning they have a backborend Spinal cord.
The study of Protochordates (first chardates
Sea Squirt-like mollusk they have an Incurrent and excurrent sighon
Lancelet-a tiny, transparent fishlike animal that lives half buried in the sandwith its head sticking out to filter plankton from the water
Acorn worm-resembles an invertebrate but has a dorsal neve and gill
These organisms and the study of them have helped to show an evolutionary link between <u>In Vertebrate</u> and <u>Vertebrate</u>
auless Fish-the first Fish to evolve
There are only 2 types still alive today, the scalampregand haghsh
They both live as <u>para Sites</u>
Lamprey inhabit <u>Cshuaries</u> from Maine to Florida. They use their <u>Sucking</u> disk on its mouth to <u>attach</u> to <u>host</u> fish. It feeds by using its <u>teethand rasping</u> on all to make a <u>hole</u> then suck out <u>blood</u> and <u>tissue</u> .
Jagfish use sharp teeth in their round mouth to burrow holes into the bodies of dead or dying fish

Cartilagrous FISh are fish whose entire SKC/Chop is composed of Cartilage
Cactilage-flexible connectivessue composed of CCILS and proteins
This and class includes Sharks, Skates, and rays
There are 700 species of <u>Cartilagino</u> Ush
Cart. fish were the first <u>Jawed</u> fish in the ocean
They have placold scales which are tiny teth that are deeply embedding 5Kin
They have visible all Slits for breathing, and also have breathing holes called spiracles located behind the lucs
\cdot
The fins of cart. fish are more Y 1910 than bony fish
SKatsand Rays
Their SCCTOXCO fins are more highly developed than Sharks
Manda Ray-are found in the MODIC and can grow to 7 meters (over 21 ft) across.
They Open their enormous MULL while Sulmming Filter feed on planktor
Strangage found, often well hidden in the Sandof the Gulf of Mexico and along the
Atlantic coast from the Carolinas to Brazil. There is a <u>Sharp Spire</u> located at the <u>base</u> of
the tall and can inflict a painful purcture, wound.
SKates - do not have a Spine on their tail and are found in temperatures of
the Atlantic and Pacific coasts.
Both Skates and Stingrays at Crustaceans and Mollusks that live in the Subtidal zone.

But to a contract

Sharks have a full cartilaginous skeleton and a highly streamlined body.
The earliest known sharks were present about 450 million years ago.
- modern sharks have been around for the past 100 million years
There are currently 440 species alive today - smallst the dwarf lantanshak Icm. or 4.7 in. long - largest - the whiteshark, 15 m or 49sh ft
Shark life-spans vary by species but most live <u>20</u> to <u>30</u> years. A few live to <u>100</u> years.
- are embedded in the Gums rather than directly affixed to the faw and are constantly replaced throughout life.
- rowsf replacementeeth grow on the Inside of the Jawand steadily move forwardike a "Canveyor belt"
- some sharks lose 30,000 or more teeth in their lifetime.
- tooth <u>Shape</u> depends on <u>diet</u> : - if they eat <u>mollasks</u> and <u>Crustacian</u> ey have <u>dense flattened</u> teeth for <u>Crushing</u>
- those that feed on Fish have needle - like teeth for gripping
- those that feed on <u>larger</u> prey have <u>pointed</u> lower teeth for <u>gripping</u> and <u>triangular</u> upper teeth with <u>Scrated edges</u> for <u>Clutting</u> .

Most sharks are " <u>COld-blooded</u> " which means that their <u>International body Jemperature</u> matches that of their <u>ambient</u> environment.
Some sharks, such as the Shortfin make and the great white maintain a higher body than the surrounding water. - this allows faster bursts of speed and more unagy
Most sharks are <u>Carnivorous</u> .
Some species, including <u>Hack</u> sharks, eat <u>Almost anything</u> out the vast <u>majority</u> seek <u>particular</u> prey, and rarely <u>Vary</u> their diet.
Sharks and buoyancy - Sharks have an enlarged <u>IVUX</u> that is filled with <u>OIJ</u> that comprises <u>30</u> % of their body weight. - <u>Carfilagis</u> about half as <u>dense</u> as <u>bone</u> , which also helps their buoyancy - Sharks also use <u>dynamic lift</u> , which means <u>Swimming</u> will keep them from <u>Sinking</u> - Sharks cannot <u>Sleep</u> very long, if at all, without <u>Sinking</u>
Shark digestion - Can take a <u>Very long</u> time
- <u>Unwanted</u> items may <u>sast</u> get past the <u>Stomach</u> , and instead the shark either <u>Vonnted</u> out and <u>CCCTS</u> unwanted items from its
mouth.
Shark Intelligence
- recent studies have indicated that many species possess <u>powerful</u> <u>problems</u> <u>Solving</u> skills, <u>Social</u> skills and <u>Curiosity</u> .

Range and habitat		
- Sharks are found in <u>all</u>	Scas They generally do not live in _	Feshwate/with a few exceptions

- Sharks are common down to depths of 2,00 6 meters (6,000 ft), and some live even deeper

Shark attacks

- The average number of <u>fatalities</u> Idwide per year between 2001 and 2006 from <u>unprovoked</u> shark attacks is <u>4.3</u>.
- Only a few sharks are dangerous to humans.

such as the bulkhark and the river shark.

- Out of more than <u>440</u> species, only <u>4</u> have been involved in a significant number of fatal, unprovoked attacks on humans: the <u>Great White</u>, <u>Oceanic Whitethy</u> <u>hgcr</u>, and <u>bwl</u> + sharks

Sharks are highly important to the marine ecosystem.

- Some are 100 predators, hunting and eating 5maller and Weaker prey which allows only the 5maller organisms to 5wvve.

- Others are bottom - feeders and act as garbagemen by cleaning up dead and deauing organisms

